The Georgia Gardener Im

www.walterreeves.com

ZOYSIAGRASS LAWN CALENDAR

MOWING HEIGHT:

1/2" - 1 1/2"; raise 1/2" higher during summer heat. Remove no more than 1/3 total height at one time.

WATER:

1 inch per week if no rainfall

FERTILIZATION:

Use any turf fertilizer twice per year. Follow label directions and rate.

IDEAL pH RANGE: 5.5-6.5

Use dolomitic lime per soil test recommendations. Can be applied at any time.

AERATION:

Use a core aerator during active growth season.

DETHATCH: If thatch exceeds .5" deep use a vertical mower with blades 1" apart; go over the lawn only two directions. Topdressing with .25" soil is most effective.

SEEDING - NEW LAWN:

1 - 2 lb. per 1000 square feet

OVERSEEDING - ESTABLISHED LAWN:

Overseeding zoysia with ryegrass is not recommended

SODDING

500 sq. ft. per pallet

WEED CONTROL:

Read product label carefully to determine which weeds are controlled and on which grasses the product can be used.

Spring pre-emergent* (prevents crabgrass, goosegrass and other summer weed seeds from sprouting)

Broadleaf post-emergent * (spot spray to kill broadleafed plants like chickweed, wild violet, dandelion, wild onion, etc)

Grassy weed post-emergent* (kills grassy weeds like crabgrass, dallisgrass, etc. Do not apply to drought-stressed Zoysia.)

Winter pre-emergent* (prevents chickweed and other winter weed seeds from sprouting)

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			ОК								
			OK								
			Possible	Best			Best	Possible			
			Possible	Best	Possible	Possible	Possible				
				Possible	Possible	Possible	Possible				
				Best	Best	Possible					
			Possible	Best	Best	Best	Possible				
	Best	Best	Marginal								
ОК	OK	ОК	OK	ОК	OK	OK	OK	OK	OK	OK	OK
				Best	Best	Possible	Possible				
								Best	Possible		

^{*} Read weed control product labels carefully. Some products can not be applied to lawns that will be seeded within a few weeks or when transitioning to active growth.